



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 5/027, 5/078, 5/083	A1	(11) International Publication Number: WO 96/33211 (43) International Publication Date: 24 October 1996 (24.10.96)
(21) International Application Number: PCT/GB96/00942 (22) International Filing Date: 22 April 1996 (22.04.96) (30) Priority Data: 9508195.6 20 April 1995 (20.04.95) GB (71) Applicants (for all designated States except US): UNIVERSITY OF BRITISH COLUMBIA [CA/CA]; Vancouver, British Columbia V6T 1Z4 (CA). UNIVERSITY OF ALBERTA [CA/CA]; Edmonton, Alberta T6G 2H7 (CA). (71) Applicant (for MN only): APPLIEDENE LIMITED [GB/GB]; 15 Clare Road, Halifax, West Yorkshire HX1 2HY (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): ANDERSEN, Raymond [CA/CA]; University of British Columbia, Dept. of Chemistry, Vancouver, British Columbia V6T 1Z4 (CA). COLEMAN, John [CA/CA]; University of British Columbia, Dept. of Chemistry & Oceanography, Vancouver, British Columbia V6T 1Z4 (CA). DE SILVA, Dilip [CA/CA]; University of British Columbia, Dept. of Chemistry, Vancouver, British Columbia V6T 1Z4 (CA). KONG, Fangming [CA/CA]; University of Columbia, Dept. of Chemistry, Vancouver, British Columbia V6T 1Z4 (CA). PIERS, Edward [CA/CA]; Uni-		versity of British Columbia, Dept. of Chemistry, Vancouver, British Columbia V6T 1Z4 (CA). WALLACE, Debra [CA/CA]; University of British Columbia, Dept. of Chemistry, Vancouver, British Columbia V6T 1Z4 (CA). ROBERGE, Michel [CA/CA]; University of British Columbia, Dept. of Biochemistry, Vancouver, British Columbia V6T 1Z4 (CA). ALLEN, Theresa [CA/CA]; 11223 72nd Avenue, Edmonton, Alberta T6G 0B4 (CA). (74) Agents: BRIERLEY, Anthony, Paul et al.; Appleyard Lees, 15 Clare Road, Halifax HX1 2HY (GB). (81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: BIOLOGICALLY ACTIVE PEPTIDES AND COMPOSITIONS, THEIR USE		
<div style="text-align: center;"> <p style="text-align: right;">(I)</p> </div>		
(57) Abstract <p>Compounds of general formula (I) wherein: R₁ and R₇₀ independently represent a hydrogen atom or an optionally substituted alkyl or acyl group; R₂ represents a hydrogen atom or an optionally substituted alkyl or acyl group or is absent when R₆ represents a group -CH= as hereinafter described; R₇₃ represents a hydrogen atom or an optional substituent or is absent when R₆ represents a methylene group or a group -CH= as hereinafter described; Y represents an optional substituent; n represents 0, 1, 2, 3, or 4; R₃ represents a hydrogen atom, or an optionally substituted alkyl group; R₇₄ represents a hydrogen atom, a hydroxy group or an optionally substituted alkyl or acyl group; R₇ represents a hydrogen atom or an alkyl group; R₇₅ represents an optionally substituted alkyl group; and i) R₆ and R₇₁ independently represent a hydrogen atom or an optionally substituted alkyl or acyl group; and R₇₂ represents a hydrogen atom; or ii) R₇₁ represents a hydrogen atom or an optionally substituted alkyl or acyl group and R₇₂ represents a hydrogen atom or R₇₁ and R₇₂ each represent radicals so that a double bond is formed between the carbon atoms to which they are attached; and R₆ represents an optionally substituted methylene group bonded to the indole moiety thereby to form a tricyclic moiety; or R₆ represents an optionally substituted group -CH= bonded to the indole moiety thereby to form an aromatic tricyclic moiety; for use in therapy and as antimitotic reagents are described. Novel hemiasterlins, criamides and geodiamolides are also described.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

BIOLOGICALLY ACTIVE PEPTIDES AND COMPOSITIONS, THEIR USE

This invention relates to novel biologically active
5 compounds and compositions, their use and derivation.

The invention has involved the extraction of novel
biologically active compounds from marine sponges.

10 Except where otherwise stated, throughout this
specification, any alkyl moiety suitably has up to 8,
especially up to 6, most preferably up to 4, carbon atoms
and may be of straight chain or, where possible, of
branched chain structure. Generally, methyl is a
15 preferred alkyl group. Halogen atoms may be fluorine,
chlorine, bromine or iodine. A preferred acyl group is
alkylcarbonyl, especially acetyl.

20 Except where otherwise stated in this specification,
optional substituents of an alkyl group may include
halogen atoms, for example fluorine, chlorine, bromine and
iodine atoms, and nitro, cyano, alkoxy, hydroxy, amino,
alkylamino, sulphinyl, alkylsulphinyl, sulphonyl,
alkylsulphonyl, amido, alkylamido, alkoxycarbonyl,
25 haloalkoxycarbonyl and haloalkyl groups. Preferably,
optionally substituted alkyl groups are unsubstituted.


30 Except where otherwise stated, throughout this
specification the recitation of a compound denotes all
possible isomers possible within the structural formula
given for those compounds, in particular geometrical and
optical isomers. Unless otherwise stated definitions are
to be regarded as covering mixtures of isomers, and
individual isomers, including racemic mixtures, where they
35 can be resolved.

- 2 -

Except if otherwise stated, definitions of compounds in this specification are to be regarded as covering all possible salts of the compounds.

5 Compounds according to the first aspect of the invention have been found to be surprisingly effective in the in vivo treatment of cancer.

10 In accordance with the first aspect of the present invention there is provided the use of a hemiasterlin compound of general formula



20 wherein:

R_1 and R_{70} independently represent a hydrogen atom or an optionally substituted alkyl or acyl group;

R_2 represents a hydrogen atom or an optionally substituted alkyl or acyl group or is absent when R_6 represents a group

25 $-CH=$ as hereinafter described;

R_{73} represents a hydrogen atom or an optional substituent or is absent when R_6 represents a methylene group or a group $-CH=$ as hereinafter described;

Y represents an optional substituent;

30 n represents 0, 1, 2, 3, or 4;

R_3 represents a hydrogen atom, or an optionally substituted alkyl group;

R_{74} represents a hydrogen atom, a hydroxy group or an optionally substituted alkyl or acyl group;

35 R_7 represents a hydrogen atom or an alkyl group;

- 3 -

R_{73} represents an optionally substituted alkyl group; and
i) R_6 and R_{71} independently represent a hydrogen atom or an optionally substituted alkyl or acyl group; and
 R_{72} represents a hydrogen atom;

5 or

ii) R_{71} represents a hydrogen atom or an optionally substituted alkyl or acyl group and R_{72} represents a hydrogen atom or R_{71} and R_{72} each represent radicals so that a double bond is formed between the carbon atoms to which they are attached; and

10 R_6 represents an optionally substituted methylene group bonded to the indole moiety thereby to form a tricyclic moiety; or

R_6 represents an optionally substituted group $-CH=$ bonded to the indole moiety thereby to form an aromatic tricyclic moiety;

15 for the manufacture of a medicament for use in therapy.

Preferably, R^1 represents a hydrogen atom or an alkyl group, especially a methyl group. More preferably, R_1 represents a hydrogen atom.

Suitably, R_2 represents a hydrogen atom or an acyl group. An acyl group may be a benzoyl group, but is preferably an alkylcarbonyl group. An especially preferred acyl group is an acetyl group. Preferably, R_2 represents a hydrogen atom.

Preferably, R_{70} represents a hydrogen atom or an alkyl group, especially a methyl group.

Where Y and/or R_{73} represent optional substituents, said substituents may be independently selected from halogen, especially fluorine, chlorine, bromine and iodine atoms and alkyl, acyl, $-OH$, $-O$ -alkyl, O -acyl, $-NH_2$, $-NH-$

- 4 -

alkyl, -N(alkyl)₂, -NH-acyl, -NO₂, -SH, -S-alkyl and -S-acyl, wherein the alkyl and acyl groups of the substituents are optionally substituted.

- 5 Preferred optional substituents represented by Y and/or R₇₃ are alkyl groups.

Preferably, R₇₃ represents a hydrogen atom.

- 10 Preferably, n represents 0, 1 or 2. More preferably, n represents 0.

Suitably, R₃ represents an alkyl group. Preferably, R₃ represents a C₃₋₆, especially C₃₋₄, branched alkyl group,
15 for example tertiary butyl or isopropyl.

Suitably, R₇₄ represents a hydrogen atom or methyl group, especially a hydrogen atom.

- 20 Suitably, R₇ represents an alkyl, preferably methyl, group.

Preferably, R₆ represents a hydrogen atom, or an optionally substituted alkyl group, or a methylene group
25 bonded to the indole moiety thereby to form a tricyclic moiety. More preferably, R₆ represents an alkyl group.


Preferably, R₇₁ independently represents a hydrogen atom or an optionally substituted alkyl or acyl group.
30 More preferably, R₇₁ represents an alkyl, especially a methyl group.

Preferably, R₆ and R₇₁ are as described in i) above.

- 35 R₇₅ may represent a group of general formula

- 5 -

5



wherein Q represents an optionally substituted $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CHCH}-$, $-\text{CH}_2\text{C.C.}-$ or phenylene moiety; and

10

X represents a group $-\text{OR}_4$, $-\text{SR}_4$, or $-\text{NR}_4\text{R}_{10}$ wherein R_4 , R_9 and R_{10} independently represent a hydrogen atom or an optionally substituted alkyl group.

15

Where Q represents one of the aforesaid optionally substituted acyclic moieties, the moiety may be substituted by one or more alkyl groups. A phenylene moiety may be substituted by one or more substituents Y as described above.

20

Where X represents a group $-\text{OR}_4$, suitably R_4 represents a hydrogen atom or a methyl group. Preferably, R_4 represents a hydrogen atom.

25

Where X represents a group $-\text{NR}_4\text{R}_{10}$, suitably R_4 represents a hydrogen atom or an alkyl group, for example a methyl group and R_{10} represents a substituted alkyl group.

30

Where R_{10} represents a substituted alkyl group, said group preferably represents a group of general formula $-\text{CHR}_{21}\text{COOH}$ wherein R_{21} represents an optionally substituted alkyl group. Preferably, R_{21} represents a group which includes at least one nitrogen atom. Preferably, R_{21}

35


represents a group of general formula $-(\text{CH}_2)_r\text{NR}_{22}\text{R}_{23}$ wherein

- 6 -

x is an integer, preferably in the range 1-4, and R_{22} and R_{23} independently represent a hydrogen atom or an optionally substituted alkyl, alkenyl or imine group. Preferably, R_{22} represents a hydrogen atom and R_{23} represents an imine group $-C(NH)NH_2$.

Preferably, X represents a group $-OR_4$, wherein R_4 represents a hydrogen atom.

Preferably, R_{75} represents a group of general formula



wherein R_4 and R_5 independently represent a hydrogen atom or an optionally substituted alkyl group; R_{76} and R_{77} each represent a hydrogen atom or a radical so that a double bond is formed between the carbon atoms to which they are attached; and X is as described above.

It has been discovered that compounds of general formula I, can cause mitotic arrest and the production of abnormal mitotic spindles. Accordingly, the invention extends, in a second aspect, to the use of a compound of general formula I as an antimitotic compound.

The compound may be used in vivo or in vitro as an antimitotic compound in, for example, procedures that require the blocking of cells in mitosis, such as the preparation of mitotic spreads for karyotype analysis and the probing of microtubule function in mitotic cells.

35

- 7 -

In a third aspect, the invention provides a novel compound of general formula I as described herein, but excluding a single compound of general formula I wherein R_1 represents methyl, R_2 represents a hydrogen, R_{70} represents methyl, R_{71} represents methyl, R_{73} represents hydrogen, n represents 0, R_3 represents t-butyl, R_{74} represents hydrogen, R_6 represents methyl, R_7 represents methyl, R_{72} represents hydrogen and R_{75} represents $-\text{CH}(\text{CH}(\text{CH}_3)_2)\text{CH}.\text{CCH}_3.\text{COOH}$. The excluded compound is hemiasterlin.

Preferably, there is provided a compound of general formula I as described herein, but excluding a compound in which R_1 represents a methyl group.

Preferably, there is provided a compound of general formula I as described herein, but excluding a compound in which R_2 represents a hydrogen atom.

One class of preferred novel compounds comprises hemiasterlins of general formula I wherein:

- R_1 represents a hydrogen atom;
- R_2 represents a hydrogen atom, or an alkyl group, or an acyl group;
- R_3 represents a hydrogen atom, or an optionally substituted alkyl group;
- n represents 0;
- R_{70} and R_{71} independently represent a hydrogen atom or optionally substituted alkyl group, but preferably each represent a methyl group;
- R_{72} , R_{73} and R_{74} represent hydrogen atoms;
- R_7 represents a hydrogen atom or an alkyl group;

- 8 -

R_4 represents a hydrogen atom, or an optionally substituted alkyl group, or a methylene group bonded to the indole moiety thereby to form a tricyclic moiety;
 R_{75} represents a group of general formula III described
5 above wherein R_4 represents a hydrogen atom, or an optionally substituted alkyl group; R_5 represents a hydrogen atom or an alkyl group; R_{76} and R_{77} represent radicals as described; and X represents a group $-OR_4$ or a group $-NR_8R_{10}$, wherein R_8 , R_9 and R_{10} independently represent
10 a hydrogen atom or an optionally substituted alkyl group.

Another class of preferred novel compounds comprises hemiasterlins of general formula I wherein:
 R_1 represents a hydrogen atom or an alkyl group;
15 R_2 represents an acyl group;
 R_3 represents a hydrogen atom, or an optionally substituted alkyl group;
 n represents 0;
 R_{70} and R_{71} independently represent a hydrogen atom or
20 optionally substituted alkyl group, but preferably each represent a methyl group;
 R_{72} , R_{73} and R_{74} represent hydrogen atoms;
 R_7 represents a hydrogen atom or an alkyl group;
 R_6 represents a hydrogen atom, or an optionally substituted
25 alkyl group, or a methylene group bonded to the indole moiety thereby to form a tricyclic moiety;
 R_{75} represents a group of general formula III described above wherein R_4 represents a hydrogen atom, or an optionally substituted alkyl group; R_5 represents a
30 hydrogen atom or an alkyl group; R_{76} and R_{77} represent radicals as described; and X represents a group $-OR_4$ or a group $-NR_8R_{10}$, wherein R_8 , R_9 and R_{10} independently represent a hydrogen atom or an optionally substituted alkyl group.

- 9 -

Another class of preferred novel compounds comprises
criamides of general formula I wherein:

R_1 represents a hydrogen atom or an alkyl group;

R_2 represents a hydrogen atom, or an alkyl group, or an
acyl group;

R_3 represents a hydrogen atom, or an optionally substituted
alkyl group;

n represents 0;

R_{70} and R_{71} independently represent a hydrogen atom or
optionally substituted alkyl group, but preferably each
represent a methyl group;


R_{72} , R_{73} and R_{74} represent hydrogen atoms;

R_6 represents a hydrogen atom, or an optionally substituted
alkyl group, or a methylene group bonded to the indole
moiety thereby to form a tricyclic moiety;

R_{75} represents a group of general formula III described
above wherein R_4 represents a hydrogen atom, or an
optionally substituted alkyl group; R_5 represents a
hydrogen atom or an alkyl group; R_{76} and R_{77} represent
radicals as described; and X represents a group $-NR_9R_{10}$,
wherein R_9 and R_{10} independently represent a hydrogen atom
or an optionally substituted alkyl group.


In formulas I and III drawn above, the bonds drawn in
wavy line are from carbon atoms which are, or may be,
optical centres.

Preferably, in the compound of general formula I the
following optical configurations predominate.



- 10 -

In accordance with a further aspect of the present invention there is provided a geodiamolide compound of general formula



wherein:

R_{31} represents an alkyl group;

R_{32} represents a hydrogen atom or an alkyl group; and


A represents a halogen atom.

Preferably, R_{31} represents a methyl group.

Suitably, R_{32} represents a hydrogen atom or, preferably, a methyl group.

Suitably, A represents a chlorine, bromine or, preferably, an iodine atom.


In the formula IV as drawn above the bonds shown in wavy line are from carbon atoms which are optical centres, except for the carbon atoms carrying moieties R_{31} and R_{32} , when those moieties are hydrogen atoms. Preferably, the following optical configuration predominates.



5
10
15
Certain compounds of the general formulae I and IV as defined above may be obtained from the marine sponge *Cymbastela* sp. (formerly classified as *Pseudaxinyssa* sp.); or be a derivatisation of compounds obtained therefrom. Derivatisation of compounds of general formula I may involve standard acylation of the extracted compounds, optionally by standard esterification. Alternatively compounds of the general formula I and IV may be prepared by entirely synthetic routes.

20
The invention extends to the use of a compound of general formula IV for the preparation of a medicament for use in therapy.

25
Compounds of general formula I may be prepared by coupling amino acid moieties A, B and C as represented below.




The coupling reactions may involve standard procedures. The amino acid moieties A, B, C may be

- 12 -


prepared by standard procedures or in procedures analogous to the procedures described in the examples hereinafter.

One general procedure for the preparation of certain
5 compounds of general formula I is provided below.


Compounds of general formula I wherein R_{75} represents



may be prepared by reacting a compound of general formula




with a compound of general formula



- 13 -

A coupling agent, for example N,N'-dicyclohexylcarbodiimide (DCC), is suitably used in the reaction. The reaction suitably comprises contacting
5 compounds V and VI in the presence of the coupling agent, a base such as triethanolamine (TEA) and an organic solvent, such as acetonitrile, suitably at a reduced temperature. After a period of time, an inorganic base, for example sodium hydroxide may be added and,
10 subsequently, the temperature raised to ambient temperature and trifluoroacetic acid (TFA) added. The desired compound of general formula I may then be isolated by standard techniques.


15 Compounds of general formula VI may be prepared by reacting a compound of general formula



25 wherein BOC (tert-butoxycarbonyl) is a protecting group, with an ylid of formula for example $(\text{Ph})_3\text{C}=\text{CR}_2\text{CO}_2\text{R}_4$. The reaction is suitably carried out in the presence of potassium carbonate in a 1:1 mixture of THF/water as described in R. Lloyd (1994) Ph.D. Thesis, University of
30 Cambridge. The protecting group BOC is suitably removed, when required, by reaction in TFA for about 2 hours at ambient temperature.

Compounds of general formula VII may be prepared by
35 reacting a compound of general formula

- 14 -




VIII

5

with a reducing agent, for example lithium aluminium hydride in tetrahydrofuran.

10 Compounds of general formula VIII may be prepared from compounds of general formula




IX

15

using the method described in Synthesis 1983, 676.

20

Compounds of general formula IX may be prepared from compounds of general formula



X

25


30

by reaction with a compound of general formula R₄I in the presence of an alkali metal hydride and in THF, using the method described in Can.J.Chem. 1973, 51, 1915.

35

- 15 -


Compounds of general formula V may be prepared by reacting a compound of general formula



10 initially with a base, for example dilute sodium hydroxide solution, followed by acidification down to about pH6. In some circumstances, it is desirable to protect the group -NR₆R₂ from reaction and this is suitably afforded using a

15 protecting agent such as BOC.

Compounds of general formula XI may be prepared by reacting a compound of general formula




with a base followed by treatment with an azide compound. The azide derivative of compound XII may then be reduced to form an amine derivative which may then be treated with

30 groups R₆₁ and/or R₂₁ in the presence of a base, for example sodium hydride, to afford the group R₆R₂N- in the compound of formula XI.

Compounds of general formula XII may be prepared by

35 coupling compounds of general formula


- 16 -



XIII

5

and



XIV


10

15

using a coupling agent, such as DCC, a base such as TEA and in an organic solvent such as acetonitrile suitably at about 0°C.

20

Compounds of general formula XIII may be prepared from a compound of general formula



XV

25

30


by processes well known to skilled persons in the art.

Compounds of general formula XIV and XV may be prepared by processes well known to skilled persons in the art.

35

- 17 -

Compounds of general formula I wherein R_7 represents



may be prepared by reacting a compound of general formula I as described above with an amine of general formula $\text{R}_9\text{R}_{10}\text{NH}$. The reaction is suitably carried out in the presence of a coupling agent such as DCC, a base such as TEA and in an organic solvent, such as acetonitrile, suitably at a reduced temperature. In certain circumstances, the group $\text{---NR}_9\text{R}_{10}$ may need protecting from undesired reactions and this is suitably afforded using a protecting agent such as BOC.

In accordance with a further aspect of the present invention there is provided a method of obtaining a compound of general formula I by extraction of a compound of general formula I from *Cymbastela* sp., including the steps of separation and purification; and optionally derivatising said compound to derive a further compound of general formula I.

Derivatisation of a compound of general formula I may include acylation and/or esterification steps. Esterification and/or acylation steps may be undertaken under standard conditions.

In accordance with a further aspect of the present invention there is provided a method of obtaining a compound of general formula IV by extraction of a compound of general formula IV from *Cymbastela* sp. including the steps of separation and purification; and optionally

- 18 -

derivatising said compound to derive a further compound of general formula IV.

5 Compounds of the general formula I and IV are biologically active. The invention further relates to the biological use of a compound of general formula I or IV. Compounds of general formula I or IV may have pesticidal, for example insecticidal activity. Preferably, however, the use is in the veterinary or, most preferably, the
10 pharmaceutical field.

 The compounds described herein may in particular have utility as antibacterial and/or antiviral agents, and/or, especially, as cytotoxic agents.

15

 The invention further provides the use of any compound of general formula I or IV for the manufacture of a medicament for use in the treatment of cancer or a tumor in a mammal.

20

 In using a compound of general formula I or IV as described in any statement herein, the compound is preferably administered to a patient in a pharmaceutical or veterinary composition comprising also a
25 pharmaceutically or veterinarily acceptable carrier, and optionally, one or more other biologically active ingredients. Such compositions may be in any form used for administering pharmaceuticals, for example any form suitable for oral, topical, vaginal, parenteral, rectal
30 and inhalatory application. The compositions may be provided in discrete dose units. The carriers may be particulate, with the compositions being, for example, tablets or powders, or liquid, with the compositions being, for example, oral syrups or injectable liquids, or
35 gaseous, for inhalatory application.

- 19 -

For oral administration an excipient and/or binder may be present. Examples are sucrose, kaolin, glycerin, starch dextrins, sodium alginate, carboxymethylcellulose and ethyl cellulose. Colouring and/or flavouring agents may be present. A coating shell may be employed. For rectal administration oleaginous bases may be employed, for example lanolin or cocoa butter. For an injectable formulation buffers, stabilisers and isotonic agents may be included.

10

The dosage of the compounds of general formula I and IV may depend upon the weight and physical condition of the patient; on the severity and longevity of the illness; and on the particular form of the active ingredient, the manner of administration and the composition employed. A daily dose of from about 0.001 to about 100 mg/kg of body weight taken singly or in separate doses of up to 6 times a day, or by continuous infusion, embraces the effective amounts most typically required. A preferred range is about 0.01 to about 50 mg/kg of body weight, per day, most preferably about 0.1 to about 30 mg/kg of body weight, per day.

It is to be understood that use of a compound of general formula I or IV in chemotherapy can involve such a compound being bound to an agent, for example a monoclonal or polyclonal antibody, a protein or a liposome, which assists the delivery of the said compound to tumour cells.

30

Therefore, the invention relates further to a pharmaceutical and veterinary composition comprising an effective amount of a compound of formula I or IV, in association with a carrier.

35

- 20 -

The invention will now be further described, by way of example with reference to Figures 1 and 2 which are graphs relating to antimitotic activity.

5 Procedure 1 : - Isolation of naturally occurring compounds

Specimens of *Cymbastela* sp. were collected by hand using SCUBA on reefs off Mandang, Papua, New Guinea. Freshly collected sponge was frozen on site and transported to Vancouver, Canada, over dry ice. The
10 sponges were identified by a leading sponge taxonomy expert Professor R. van Soest. A voucher sample was been deposited at the Institut voor Systematiek en Populatiebiologie-Zoologisch Museum, University of Amsterdam.

15

The thawed sponge (260 g dry wt) was extracted exhaustively with a solution of $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (1:1). Evaporation of the organic extract *in vacuo* gave an aqueous suspension. MeOH was added to give a 9:1 MeOH:H₂O
20 solution (1 l), which was extracted with hexanes (4 x 250 ml). The hexane extracts were combined and concentrated *in vacuo* to yield an orange oil. Water was added to the MeOH solution to give a 1:1 MeOH/H₂O solution, which was extracted with CHCl_3 (4 x 250 ml). The combined CHCl_3 ,
25 layers were concentrated *in vacuo* to yield an orange oil (3.5 g). Repeated size exclusion chromatography on Sephadex LH-20 eluting with MeOH yielded a number of crude geodiamolides and hemiasterlins. Pure geodiamolide G (Compound 1 below) (2 mg, 0.0007% dry wt) was obtained via
30 reversed-phase HPLC (MeOH/H₂O 60:40). Reversed-phase isocratic HPLC (0.05% TFA:MeOH 1:1) afforded hemiasterlin A (Compound 2 below - 32 mg, 0.012% dry wt) and hemiasterlin B (Compound 3 below - 1 mg, 0.0004% dry wt). These compounds are novel. The reversed-phase isocratic
35 HPLC using TFA and MeOH also yielded the known compound

- 21 -

hemiasterlin (Compound A below - 40 mg, 0.015% dry wt). This was used to prepare a novel acylated and esterified compound described later.

5 **Geodiamolide G (Compound 1):** colourless glass; IR (neat) 3313, 2977, 2933, 1732, 1675, 1635, 1505, 1455, 1417, 1377, 1285, 1217, 1102, 1083, 1052, 952, 827, 754 cm^{-1} ; NMR data, Table 1 below; HREIMS, M^+ m/z 655.1760 ($\text{C}_{28}\text{H}_{38}\text{N}_3\text{O}_7\text{I}$ ΔM 0.6 mmu).

10

Hemiasterlin A (Compound 2): white solid $[\alpha]_D = -45^\circ$ (c 0.25, MeOH); UV (MeOH) λ_{max} (ϵ) 218 (23,400), 280 nm (2,800); IR (neat) 3418, 2966, 1689, 1680, 1643 cm^{-1} ; NMR data, Table 2 below; HRFABMS, MH^+ m/z 513.3471 ($\text{C}_{29}\text{H}_{43}\text{O}_4\text{N}_4$ ΔM 3.0 mmu).

15

Hemiasterlin B (Compound 3): white solid; CD(MeOH) $(\theta)_{226}$ 10,800; NMR data, Table 2 below; HRFABMS, MH^+ m/z 499.3319 ($\text{C}_{28}\text{H}_{43}\text{O}_4\text{N}_4$ ΔM 3.4 mmu).

20

Hemiasterlin (Compound A): white solid; $[\alpha]_D = -77^\circ$ (c 0.07, MeOH); UV (MeOH) λ_{max} (ϵ) 216 (15,400), 273 nm (1,600); IR (neat) 3412, 2962, 1650, 1635 cm^{-1} ; NMR data, Table 2 below; HRFABMS, MH^+ m/z 527.3594 ($\text{C}_{30}\text{H}_{47}\text{O}_4\text{N}_4$ ΔM -0.35 mmu).


25

Product identification, including assignment of stereochemical configurations, was achieved by a range of techniques, including NMR, mass spectroscopy and optical rotation measurements, with cross-reference to analyses reported in the literature, for compounds already known. In the case of compounds 2 and 3 CD and chemical degradation analyses were carried out to assist the stereochemical determination.

30


35

- 22 -



Compound 1

20



Compound 2 $R_1 = R_2 = R_4 = H, R_{33} = Me$

Compound 3 $R_1 = R_2 = R_3 = R_{44} = H$

Compound A $R_1 = R_{33} = Me, R_2 = R_{44} = H$

TABLE 1

NMR Data for Geodiamolide G (Compound 1). Recorded in CDCl₃ at 500 MHz.

Carbon	$\delta^{13}\text{C}^a$	$\delta^1\text{H}$	COSY	HMBC ^b
1	174.5			H3, H3', H14, H22, NH(14)
2	41.0	2.46, m	H3, H3', H22	H3', H22
3	37.7	2.55, ddJ=12.3, 3.7Hz	H2, H3'	H22, H23, H23'
3'		2.14, tJ=11.9Hz	H2, H3	
4	143.6			H3, H23'
5	205.1			H3', H23, H23', H24
6	36.2	2.95	H7', H24	H24
7	38.8	1.82, dddJ=14.6, 9.5, 2.8Hz	H7', H8	H24, H25
7'		1.61, dddJ=14.6, 10.9, 2.8Hz	H6, H7	
8	69.7	5.11, m	H7, H25	H25
9	170.6			H10, H26
10	49.1	4.51, dqJ=7.3, 7.2Hz	H26, NH(10)	H26
NH(10)		6.35, dJ=7.3Hz	H10	
11	168.9			H12, H15'
12	57.1	5.06, ddJ=7.9, 8.9Hz	H15, H15'	H15, H15', H27
13	174.4			H12, H14, H27, H28, NH(14)
14	45.1	4.72, dqJ=7.0, 6.9Hz	H28, NH(14)	H28
NH(14)		6.19, dJ=7.0Hz	H14	
15	33.2	3.12, ddJ=14.6, 7.9Hz	H12, H15'	H12, H17, H21
15'		2.90, ddJ=14.6, 8.9Hz	H12, H15	
16	130.3			H15, H15', H20
17	138.2	7.45, dJ=1.4Hz	H21	H15, H15', H21
18	85.2			H17, H20
19	154.5			H17, H21
20	115.6	6.88, dJ=8.2Hz	H21	
21	130.6	7.04, ddJ=8.2, 1.4Hz	H17, H20	H17
22	17.9	1.15, dJ=6.4Hz	H2	
23	127.8	5.90, s		H3'
23'		5.78, s		
24	17.8	1.09, dJ=7.1Hz	H6	
25	20.8	1.28, dJ=6.3Hz	H8	
26	18.3	1.32, dJ=7.2Hz	H10	H10
27	30.7	2.97, s		H12
28	19.2	1.04, dJ=6.9Hz	H14	

^aObtained from HMQC and HMBC spectra only. ^bProton resonances that are correlated to the carbon resonance in the $\delta^{13}\text{C}$ column.

- 24 -

TABLE 2

NMR Data for the hemiasterlins, Compounds 2, 3 and A.
Recorded in DMSO-d₆ at 500 MHz.

Carbon	Hemiasterlin		Hemiasterlin A			Hemiasterlin B		
	$\delta^{13}\text{C}$	$\delta^1\text{H}$	$\delta^{13}\text{C}$	$\delta^1\text{H}$	HMBC ^a	$\delta^{13}\text{C}^b$	$\delta^1\text{H}$	HMBC ^a
1-N				10.93.s			10.88.s	
2	128.7	7.16.s	122.9	7.11.s	H1	122.9	7.06.s	
3	116.5		120.4		H2,5,6,14,15	119.0		H14,15
4	125.0		124.8		H1,2,8	124.8		H2,8
5	120.6	8.09.dJ=8 Hz	120.2	7.80.dJ=8 Hz		120.0	7.98.dJ=7.8 Hz	
6	121.1	7.07.tJ=8 Hz	120.7	7.06.tJ=8 Hz		120.4	7.05.tJ=7.8 Hz	
7	118.4	7.20.tJ=8 Hz	118.1	6.96.tJ=8 Hz	H8	117.8	6.95.tJ=7.8 Hz	H8
8	110.0	7.44.dJ=8 Hz	111.8	7.35.dJ=8 Hz		111.4	7.38.dJ=7.8 Hz	H7
9	137.7		137.3		H1,2,5,6	137.2		H2
10	37.5		37.9		H11,14,15	37.5		H14,15
11	67.5	4.44.dJ=6 Hz	71.7	3.47.s	H14,15,17	69.3	3.47.bs	
12	166.0		171.2		H11			
13	32.4	3.75.s						
14	27.0	1.41.s	27.5	1.41.s	H15	27.3	1.38.s	H15
15	22.5	1.38.s	23.2	1.37.s	H11,14	22.5	1.34.s	H14
16-N		7.38.bs						
17	33.4	2.24.s	35.2	1.92.s	H11	35.0	1.93.s	
18-N		8.87.s		7.84.bdJ=9 Hz				
19	56.2	4.84.dJ=8 Hz	53.8	4.79.dJ=9 Hz		53.8	4.58.tJ=8 Hz	H22,23
20	170.1		170.9		H19,30	171.0		H30
21	34.6		34.7		H19,22,23,24	30.0	2.11.m	H19,22
22	26.3	0.99.s	26.2	0.93.s	H19,23,24		0.84.dJ=6.3 Hz	
23	26.3	0.99.s	26.2	0.93.s	H19,22,24		0.89.dJ=6.3 Hz	
24	26.3	0.99.s	26.2	0.93.s	H19,22,23			
26	55.6	4.93.tJ=10 Hz	55.9	4.91.tJ=9 Hz	H30,32,33	55.8	4.87.tJ=10 Hz	H30,32,33
27	138.3	6.66.dJ=10 Hz	138.2	6.63.dJ=9 Hz	H26,34	137.7	6.63.dJ=8.8 Hz	H26,34
28	131.6		131.8		H26,34	131.5		H34
29	168.5		168.6		H27,34	168.5		H34
30	31.1	3.03.s	30.9	2.97.s	H26	30.0	2.98.s	
31	28.7	2.01.m	28.8	1.96.m	H26,32,33	28.7	2.08.m	H32,26
32	19.3	0.80.dJ=7 Hz	19.3	0.77.dJ=6.5 Hz	H33		0.74.dJ=6.1 Hz	
33	18.9	0.78.dJ=7 Hz	18.7	0.70.dJ=6.5 Hz	H32		0.80.dJ=6.1 Hz	
34	13.5	1.80.s	13.5	1.77.s	H27	13.0	1.75.s	H27

^aProton resonances that are correlated to the carbon resonance in the $\delta^{13}\text{C}$ column.

^bObtained from HMQC and HMBC correlations.


- 25 -

In another example, specimens of *Cymbasyela* sp. were collected by hand using SCUBA on the reefs of Madang, Papua, New Guinea. Fresh sponge was frozen on site and transported to Vancouver, Canada, over dry ice. The freeze-dried sponge (157 g dry wt.) was extracted sequentially with hexane, carbon tetrachloride, chloroform and methanol (3 x 8 litres). The extracts were concentrated *in vacuo* to yield 6 g, 0.76 g, 1.24 g and 1.1 g respectively for each solvent. Repeated size exclusion chromatography of the chloroform extract, on Sephadex LH-20 with methanol, yielded a mixture of crude hemiasterlins and criamides. Reversed phase HPLC, 50:50 0.05% TFA:MeOH afforded pure Hemiasterlin (60 mg, 0.038% dry wt.) (Compound A above); Hemiasterlin-A (55 mg, 0.035% dry wt.) (Compound 2 above); Hemiasterlin-B (3 mg, 0.0019% dry weight) (Compound 3 above); Criamide-A (2 mg, 0.0013% dry wt.) (Compound 6 below); Criamide-B (2 mg, 0.0013% dry wt.) (Compound 7 below).

20

25

30

Compound 6 $R_1 = \text{Me}$ Compound 7 $R_1 = \text{H}$

35

SUBSTITUTE SHEET (RULE 26)

- 26 -

Product identification of compounds 6 and 7, including assignment of stereochemical configurations, was achieved by a range of techniques as described above for the other examples. NMR data is provided in Table 3
5 below.

- 27 -

TABLE 3

NMR Data for Criamide B (Compound 7) and Criamide A (Compound 6)


Carbon no.	Criamide B				Criamide A	
	$\delta^{13}\text{C}$ (ppm) (125 MHz)	$\delta^1\text{H}$ (ppm) (500MHz)	COSY (500 MHz)	HMBC (500 MHz)	$\delta^1\text{H}$ (ppm) (500MHz)	COSY (500 MHz)
1-N		11.14 s				
2	124.2	7.16 d. J=2.3			7.17 s	
3	117.1			H1,2,14,15		
4	137.7			H5,7		
5	120.2	8.08 d. J=7.3	H6	H7	8.10 d. J=7.9	H6
6	118.0	7.03 t. J=7.3	H5,7	H8	7.08 t. J=7.0	H5,7
7	120.9	7.12 t. J=7.3	H6,8	H5	7.21 t. J=8.2	H6,8
8	112.0	7.41 d. J=8.0	H7	H6	7.46 d. J=8.5	H7
9	124.7			H1,2,6	-	
10	37.6			H14,15		
11	67.6	4.45 d. J=9.0	H16-A	H14,15	4.44 d. J=9.2	H16-A
12	165.8			H19		
13					3.75 s	
14	27.2	1.42 s		H15	1.42 s	
15	22.0	1.38 s		H14	1.38 s	
16		A 7.30 bs B 8.77 bs	H16-B,11,17 H16-A,17		7.35 bs 8.84 bs	H16-B,11,17 H16-A,17
17	33.5	2.23 t. J=4.8	H16A,16B		2.24 bs	H16A,16B
18-N		8.86 d. J=9.2	H19		8.87 d. J=6.1	H19
19	55.5	4.88 d. J=8.2	H18	H22,23,24	4.88 d. J=8.5	H18,H21
20	170.1			H19,26,30		
21	34.9			H22,23,24		
22	26.3	1.0 s		H19,23,24	1.00 s	
23	26.3	1.0 s		H19,23,24	1.00 s	
24	26.3	1.0 s		H19,22,23	1.00 s	
26	55.9	4.98 t. J=9.8	H27,31	H30,32,33	4.98 t. J=9.8	H27,31
27	132.4	6.30 d. J=9.0	H26	H26,34	6.30 d. J=8.5	H26
28	135.4			H26,34		
29	168.7			H27,29,34		
30	30.9	3.04 s		H26	3.01 s	
31	28.7	2.01 m	H26,32,33	H32,33	1.94 m	H26,32,33
32	19.5	0.83 d. J=6.5	H31	H33	0.83 d. J=7.7	H31
33	19.2	0.78 d. J=6.5	H31	H32	0.77 d. J=7.7	H31
34	14.2	1.82 s		H27	1.82 s	
35-N		8.02 d. J=7.6	H36		8.03 d. J=7.6	H36
36	51.8	4.27 m	H35,38-A,38-B		4.26 dd J=4.0,9.0	H35,38-A,38-B
38	27.7	A 1.81 m B 1.67 m	H36,38-B H38-A,39		A 1.79 m B 1.69 m	H36,38-B H38-A,39
39	24.6	1.51 m	H38-B,40		1.49 m	H38-B,40
40	40.0	3.10 dd J=5.9,7.1	H39,41		3.09 dd J=6.0,7.0	H39,41
41-N		7.53 t. J=5.9	H40		7.59 t. J=5.0	H40

SUBSTITUTE SHEET (RULE 26)

- 28 -

Procedure 2 : Derivatisation of naturally occurring compounds

The following compounds 4 and 5, were respectively
5 prepared from compounds A and 2 above.



Compound 4 $R_1 = R_3 = R_4 = \text{Me}$, $R_2 = \text{COCH}_3$

Compound 5 $R_1 = \text{H}$, $R_2 = \text{COCH}_3$, $R_3 = R_4 = \text{Me}$

20 They were prepared by firstly converting compounds A and 2 to their methyl esters, then by acylating the esters.

25 To prepare the methyl esters of compounds A and 2, diazomethane was prepared in the standard fashion from 1-methyl-3-nitro-1-nitrosoguanidine (MNNG) using an Aldrich MNNG-diazomethane kit. The resulting yellow diazomethane solution (3 ml ether) was added to 1 mg of peptide A or 2, dissolved in 3 ml of chloroform. The reaction mixture was left at ambient temperature for one hour then the solvents
30 were removed in *vacuo*.

To then acylate the esterified peptides, approximately 1 mg of each esterified peptide was stirred overnight under an argon atmosphere with freshly distilled
35 (from NaOH) pyridine and acetic anhydride (1.0 ml each).

- 29 -


Excess reagents were removed in vacuo to yield the pure N-acetyl peptide esters, compounds 4 and 5, which are characterised as follows.

- 5 **Hemiasterlin, N-acetyl methyl ester (Compound 4):** white solid; CD(MeOH) (θ)₂₃₁ + 13,300; ¹H NMR CD₂Cl₂, 500 MHz) δ 0.44 (s), 0.84 (d, J = 6.6 Hz), 0.87 (d, J = 6.6 Hz), 1.39 (s), 1.59 (s), 1.86 (s), 2.04 (m), 2.16 (s), 2.93 (s), 3.15 (s), 3.70 (s), 3.76 (s), 4.41 (d, J = 9 Hz), 5.03 (t, J = 9 Hz), 6.18 (d, J = 8 Hz), 6.39 (s), 6.64 (d, J = 8 Hz), 7.10 (s), 7.15 (t, J = 8 Hz), 7.24 (t, J = 8 Hz), 7.32 (d, J = 8 Hz), 8.29 (d, J = 9 Hz); EIHRMS, M⁺ m/z 582.3796 (C₃₃H₅₀N₄O₅, Δ M 1.5 mmu).
- 10
- 15 **Hemiasterlin A, N-acetyl methyl ester (Compound 5):** white solid; CD(MeOH) (θ)₂₃₁ +10,400; ¹H NMR CD₂Cl₂, 500 MHz) δ 0.48 (s), 0.83 (d, J = 6.6 Hz), 0.88 (d, J = 6.6 Hz), 1.40 (s), 1.54 (s), 1.85 (s), 1.98 (m), 2.16 (s), 2.93 (s), 3.15 (s), 3.70 (s), 4.48 (d, J = 10.1 Hz), 5.02 (t, J = 10 Hz), 6.19 (d, J = 9 Hz), 6.37 (s), 6.65 (d, J = 10.5 Hz), 7.17 (t, J = 7 Hz), 7.20 (t, J = 7 Hz), 7.23 (s), 7.40 (d, J = 7 Hz), 8.30 (d, J = 10 Hz), 8.31 (s); EIHRMS, M⁺ m/z 568.3626 (C₃₂H₄₈N₄O₅, Δ M 0.1 mmu).
- 20

- 30 -

Procedure 3 : Totally synthetic method

The process for preparing compounds described herein by a totally synthetic method involves, in general terms, the coupling of amino acids. Thus, the preparation of the compound



involves coupling amino acids corresponding to units A, B and C. Triamide compounds which include an additional moiety D coupled to end C of the compound shown above can be prepared by coupling an amino acid corresponding to the desired moiety D to the peptide A-B-C.

20 The preparations of amino acids A and C are described below. Amino acid B is commercially available.

1. Preparation of amino acid C (N-methylhomo vinyllogous valine ethyl ester)

25

Scheme 1 below provides a summary of the procedures described hereinafter.

(a) **Preparation of N-Boc,N-Me-L-valine (1):**

30

N-Boc-L-valine (5 g; 23 mmol) and methyl iodide (1.59 ml; 25.3 mmol) were dissolved in THF (65 ml) and the solution was cooled to 0°C under argon. Sodium hydride dispersion (2.03 g; 50.6 mmol) was added cautiously with gentle stirring. The suspension was stirred at room

- 31 -

temperature for 16 h. Ethyl acetate (30 ml) was then added (to consume the sodium hydroxide formed from the excess sodium hydride), followed by water, drop wise, to destroy the excess sodium hydride. The solution was
5 evaporated to dryness, and the oily residue partitioned between ether (25 ml) and water (50 ml). The ether layer was washed with 5% aqueous NaHCO₃ (25 ml), and the combined aqueous extracts acidified to pH 3 with aqueous citric acid. The product was extracted into ethyl acetate (3 x
10 25 ml), the extract washed with water (2 x 25 ml), 5% aqueous sodium thiosulfate (2 x 25 ml; to remove iodine), and water (2 x 25 ml), dried over MgSO₄ and evaporated to give a yellow oil. The procedure was repeated to improve the overall yield. Final yield was 3.53 g; 70.6%.

15 ¹H nmr (CDCl₃; 400 MHz) δ 0.86 (d, 3H, J=7 Hz), 0.97 (d, 3H, J=7 Hz), 1.41 (s, 9H), 2.17 (bs, 1H), 2.80 & 2.82 (2 s, 3H), 4.05 (d, 0.5H, J=10Hz), 4.26 (d, 0.5H, J=10Hz), 10.8 (bs, 1H).

20 (b) Preparation of N-Boc,N-Me-L-valine-N-Me,N-Ome
(Weinreb amide) (2):

N-Boc,N-Me-L-valine (3.2 g; 13.9 mmol), N,O-dimethylhydroxylamine hydrochloride (1.5 g; 15 mmol) and
25 triethyl amine (2.1 ml; 30 mmol) were dissolved in CH₂Cl₂ (30 ml) and the solution cooled to -10°C under argon. Dicyclohexylcarbimide (3.1 g; 15 mmol) was dissolved in 15 ml of CH₂Cl₂ and added drop wise to the reaction mixture over 10 minutes. The solution was stirred for an
30 additional 15 minutes at -10°C and then for 1 h at room temperature at which time it was filtered and the excess solvents removed in vacuo. The oil was dissolved in EtOAc (50 ml) and washed with 5% HCl (2 x 25 ml), water (2 x 25 ml), 5% NaHCO₃ (2 x 25 ml), and water (2 x 25 ml), dried

- 32 -

over MgSO_4 and evaporated to yield a yellow oil (3.4 g; 85% yield).

^1H nmr (CDCl_3 ; 400 MHz) δ (0.82 (t, 6H $J = 7$ Hz), 1.38 (s, -3 H), 1.41 (s, -6H), 2.16 (m, 1H), 2.73 (s, -1H), 2.76 (s, -2H), 3.13 (s, 3H), 3.62 (s, -1H), 3.65 (s, -2H). Doubling of peaks caused by rotamers around the N-Boc group.

(c) Preparation of N-Boc, N-Me-L-valine aldehyde (3):

10

Weinreb amide (226 mg; 0.78 mmol) was dissolved in THF (8 ml) and cooled to -78°C , then added drop wise to a dispersion of LiAlH_4 (35 mg; 0.86 mmol) in THF at -78°C . The reaction was stirred for 0.5 h at which time it was quenched with $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$ (251 mg; 0.78 mmol) and allowed to warm to room temperature. The solution was filtered through celite and the excess solvents removed in vacuo to yield a colourless oil. Normal phase silica gel chromatography, eluting with 1:6 ethyl acetate: hexanes, afforded pure N-Boc, N-Me-L-valine aldehyde as a clear oil (116 mg; 68%).

^1H nmr (CDCl_3 ; 400 MHz) δ 0.84 (d, 3H, $J=7$ Hz), 1.03 (d, 3H, $J=7$ Hz), 1.40 (s, 9H), 2.20 (bs, 1H), 2.72 (s, 2H), 2.84 (s, 1H), 3.55 (d, 0.5H, $J=10\text{Hz}$), 4.02 (d, 0.5H, $J=10\text{Hz}$), 9.58 (bs, 1H).

(d) Preparation of N-Boc-MHVV-OEt (4):

N-Boc, N-Me-valine aldehyde (120 mg; 0.56 mmol) was dissolved in a 1:1 mixture of THF: H_2O (6 ml) and the (carbethoxy ethylidene)triphenylphosphane (222 mg; 6.2 mmol) was added and the reaction stirred at room temperature for 4 h. Normal phase silica gel chromatography, eluting with 1:6 ethyl acetate: hexanes,

- 33 -

afforded the N-Boc-homo vinylogous valine ethyl ester as a clear oil (212 mg; 71%).

¹H nmr (CDCl₃; 400 MHz) δ 0.82 (d, 3H, J=7 Hz), 0.87 (d, 3H, J=7 Hz), 1.27 (t, 3H, J = 7 Hz), 1.42 (s, 9H), 1.85 (m, 1H), 1.87 (s, 3H), 2.69 (bs, 3H) 4.17 (q, 2H, 7 Hz), 4.28 (bs, 0.5H), 4.53 (bs, 0.5H), 6.62 (d, 1H, J=9 Hz).

(e) Preparation of MHVV-OEt (5):

10

N-Boc-MHVV-OEt (200 mg; 67 mmol) was dissolved in 1 ml of 1:1 CH₂Cl₂: trifluoroacetic acid mixture and stirred under argon for 0.5 h. Excess solvents were removed in vacuo and the oily residue was twice redissolved in CH₂Cl₂ (25 ml) and concentrated to remove any traces of TFA. The final product was a white amorphous solid (207 mg, 99%).

¹H nmr (CDCl₃; 400 MHz) δ 0.82 (t, 6H, J = 7 Hz), 1.27 (t, 3H, J = 7 Hz), 1.42 (s, 9H), 1.85 (m, 1H), 1.87 (s, 3H), 2.69 (bs, 3H), 3.9 (m, 1H), 4.17 (q, 2H, 7 Hz), 6.62 (d, 1H, J = 9 Hz) 8.1 (bs, 0.5H), 8.3 (bs, 0.5H), 12.9 (bs, 1H).

2. Preparation of amino-acid A (N-Boc-tetramethyltryptophan derivative)

25

Scheme 2 below provides a summary of the procedures described hereinafter for the preparation of amino acid A and derivatives which are described herein:

30

(a) Preparation of Methyl ester (7)

To a stirred suspension of indol-3-acetic acid (6) (1.07 g, 6.11 mmol) in ether (20 ml) at room temperature was added an ethereal solution of diazomethane drop wise

35

- 34 -

until the yellow colour of the diazomethane persisted in the reaction mixture, and tlc analysis showed complete consumption of starting material. Excess diazomethane was removed under a stream of argon and the remaining solvent removed in vacuo. The crude oil thus obtained was purified by flash column chromatography (50% ether in pet. ether), to afford methyl ester (7) as an off-white solid (1.16 g, 100%).

10 Mp: 47-48°C

IR (CHCl₃ soln): 3409 (s, NH), 1729 (s, C=O), 1621 (w, C=C).

15 ¹H NMR (400 MHz, CDCl₃) δ: 3.71 (3H, s, OCH₃), 3.79 (2H, s, CH₂CO₂CH₃), 7.03 (1H, s, =CHNH), 7.15 (1H, t, J=7.8 Hz, ArH), 7.20 (1H, t, J=7.8 Hz, ArH), 7.31 (1H, d, J=7.8 Hz, ArH), 7.63 (1H, d, J=7.8 Hz, ArH), 8.12 (1H, br, s, NH).

20 ¹³C NMR (75.3 MHz, CDCl₃) δ : 172.6, 136.0, 127.1, 123.1, 122.1, 119.6, 118.7, 111.2, 108.1, 51.9, 31.9.

LRMS (EI): 189 (M⁺, 25%), 130 (100%).

25 HRMS (EI) Calcd. for C₁₁H₁₁O₂N: 189.07898; found : 189.07866.

Anal. Calcd. for : C₁₁H₁₁O₂N: C, 69.83; H, 5.86; N, 7.40.
Found: C, 69.47; H, 5.91; N, 7.50

30 (b) Preparation of Di methyl ester (8)

To a stirred, cooled (-78°C) suspension of potassium bis(trimethylsilyl) amide (4.90 g 24.6 mmol) in dry THF (100 ml) under argon was added a solution of methyl ester (7) (1.57 g, 8.31 mmol) in THF (30 ml + 20 ml washings)

- 35 -

via cannula. The reaction mixture was warmed to 0°C and stirred for two hours before recoiling to -78°C. Freshly distilled methyl iodide (5.2 ml, 82.8 mmol) was added, the mixture allowed to warm to 0°C and stirring continued for
5 three hours or until tlc analysis showed complete reaction. The reaction was quenched with water (100 ml) and then extracted with ether (3 x 100 ml), the combined organic extracts were washed with brine (100 ml), dried with magnesium sulfate and concentrated *in vacuo*. The
10 resulting crude oil was purified by flash column chromatography (25% ether in pet.ether) to afford methyl ester (8) as a viscous pale yellow oil (1.61 g, 89%).

IR (neat): 1734(s, C=O), 1615, 1550 (w, C=C).

15

¹H NMR (400 MHz, CDCl₃) δ: 1.61 (3H, d, J=7.1 Hz, CH(CH₃)), 3.67 (3H, s, NCH₃), 3.75 (3H, s, OCH₃), 4.01 (1H, q, J=7.1 Hz, CH(CH₃)CO₂CH₃), 7.00 (1H, s=CHNCH₃), 7.13 (1H, t, J=7.8 Hz, ArH), 7.24 (1H, t, J=7.8 Hz, ArH), 7.30 (1H, d, J=7.8 Hz, ArH),
20 7.68 (1H, d, J=7.8 Hz, ArH).

¹³C NMR (75.3 MHz, CDCl₃) δ: 175.6, 136.9, 126.7, 126.2, 121.7, 119.2, 119.0, 113.9, 109.2, 51.9, 36.7, 32.7, 18.0.

25 LRMS (EI): 217 (M⁺, 18%), 158 (100%).

HRMS (EI) Calcd. for C₁₃H₁₅O₂N: 217.11028; found: 217.11013.

Anal. Calcd. for : C₁₃H₁₅O₂N: C, 71.87; H, 6.96; N, 6.45.

30 Found: C, 71.52; H, 6.80; N, 6.26.

(c) Preparation of Tri methyl ester (9)

To a stirred, cooled (-78°C) suspension of potassium
35 bis(trimethylsilyl)amide (2.90 g, 14.5 mmol) in dry THF

- 36 -

(60 ml) under argon was added a solution of methyl ester (8) (1.25 g, 5.76 mmol) in THF (30 ml + 20 ml washings) via cannula. The reaction mixture was warmed to 0°C and stirred for two hours before re-coiling to -78°C. Freshly distilled methyl iodide (3.5 ml, 57.6 mmol), was added, the mixture allowed to warm to 0°C and stirring continued for three hours or until tlc analysis showed complete reaction. The reaction was quenched with water (60 ml) and then extracted with ether (3 x 75 ml), the combined organic extracts were washed with brine (75 ml), dried with magnesium sulfate and concentrated *in vacuo*. The resulting crude oil was purified by flash column chromatography (20% ether in pet.ether) to afford the methyl ester (9) as an off-white solid (1.22 g 92%).

Mp: 99-101°C

IR (CHCl₃, soln): 1727 (s, C=O), 1618, 1550 (w, C=C).

¹H NMR (400 MHz, CDCl₃) δ: 1.70 (6H, s, CCH₃(CH₃)), 3.64 (3H, s, NCH₃), 3.75 (3H, s, OCH₃), 6.94 (1H, s = CHNH), 7.10 (1H, t, J=7.9 Hz, ArH), 7.22 (1H, t, J=7.9 Hz, ArH), 7.29 (1H, d, J=7.8 Hz, ArH), 7.64 (1H, d, J=7.8 Hz, ArH).

¹³C NMR (75.3 MHz, CDCl₃) δ: 177.6, 137.4, 125.9, 125.2, 121.5, 120.2, 119.1, 119.0, 109.3, 52.1, 41.9, 32.7, 26.3.

LRMS (EI): 231 (M⁺, 15%), 172 (100%).

HRMS (EI) Calcd. for C₁₄H₁₇O₂N: 231.12593; found: 231.12572.

Anal. Calcd for : C₁₄H₁₇O₂N: C, 72.70; H, 7.41; N, 6.06.
Found: C, 72.83; H, 7.44; N, 6.04.

(d) Preparation of Alcohol (10)

- 37 -

To a stirred cooled (- 78°C) solution of methyl ester (9) (1.38 g, 5.97 mmol) in dry ether (70 ml) under an argon atmosphere was added DIBAL (14.9 ml, 1.0M in hexanes, 14.9 mmol). The resulting solution was allowed to warm to 0°C and stirring was continued for three hours. The reaction was quenched by addition of water (30 ml), allowed to warm to room temperature whereupon Rochelles salt (30 ml) was added. The organic layer was separated and the aqueous layer was extracted with ether (2 x 50 ml). The combined organic extracts were washed with brine (50 ml), dried with magnesium sulfate and concentrated in vacuo. The crude mixture was purified by flash column chromatography (50% ether in pet.ether) to afford the alcohol (10) as a white solid (1.14 g, 94%).

15

Mp: 80-82°C.

IR (CHCl₃ soln): 3400 (br, s, OH), 1614, 1545 (w, C=C).

¹H NMR (400 MHz, CDCl₃) δ: 1.48 (6H, s, CCH₃(CH₃)), 3.75 (3H, s, NCH₃), 3.79 (2H, s, CH₂OH), 6.90 (1H, s = CHNH), 7.11 (1H, t, J=7.9 Hz, ArH), 7.22 (1H, t, J=7.9 Hz, ArH), 7.32 (1H, d, J=7.8 Hz, ArH), 7.78 (1H, d, J=7.8 Hz, ArH).

¹³C NMR (75.3 MHz, CDCl₃) δ: 137.9, 127.1, 126.1, 121.5, 121.0, 119.4, 118.8, 109.6, 71.6, 37.7, 32.7, 25.5.

LRMS (EI): 203 (M⁺, 17%), 172 (100%).

HRMS (EI) Calcd. for C₁₃H₁₇ON: 203.13101; found: 203.13052.

Anal. Calcd for : C₁₃H₁₇ON: C, 76.81; H, 8.43; N, 6.89.
Found: C, 76.89; H, 8.43; N, 6.70.

(e) Preparation of Aldehyde (11)

- 38 -

To a mixture of alcohol (10) (362 mg, 1.78 mmol) 4-methylmorpholine N-oxide (375 mg, 3.21 mmol) and 4A powdered molecular sieves (400 mg) in dry dichloromethane (12 ml) under an argon atmosphere at room temperature was added solid TPAP (35 mg, 0.0996 mmol) in one portion. The resulting black mixture was stirred at the same temperature for three hours, then filtered through celite to remove the molecular sieves and the filtrate concentrated in vacuo. The black oil was purified by flash column chromatography (20% ether in pet.ether) to afford the aldehyde (11) as an off-white solid (314 mg, 88%).

Mp: 61-63°C

IR (CHCl₃ soln): 1718 (s, C=O), 1610, 1542 (w, C=C).

¹H NMR (400 MHz, CDCl₃) δ: 1.58 (6H, s, CCH₃(CH₃)), 3.70 (3H, s, NCH₃), 6.96 (1H, s, =CHNH), 7.10 (1H, dt, J= 0.9, 8.0 Hz, ArH), 7.24 (1H, dt, J=0.9, 8.0 Hz ArH), 7.32 (1H, dd, J=0.9, 8.0 Hz, ArH), 7.56 (1H, dd, J=0.9, 8.0 Hz, ArH), 9.49 (1H, s, CHO).

¹³C NMR (75.3 MHz, CDCl₃) δ: 202.2, 137.6, 126.6, 126.1, 121.8, 120.1, 119.3, 115.0, 109.5, 46.5, 32.8, 21.9.

LRMS (EI): 201 (M⁺, 13%), 172 (100%).

HRMS (EI) Calcd. for C₁₃H₁₃ON: 201.11537; found: 201.11473.

Anal. Calcd for : C₁₃H₁₃ON: C, 77.58; H, 7.51; N, 6.96. Found: C, 77.42; H, 7.58; N, 6.83.

(f) Preparation of Enol ether (12)

- 39 -

To a stirred suspension of methoxymethyltriphenylphosphonium chloride (3.03 g, 8.84 mmol) in dry THF (40 ml) under an argon atmosphere at room temperature (water bath) was added potassium tert-butoxide (991 mg, 8.82 mmol) as a solid in one portion. The reaction mixture immediately turned a deep red colour, the water bath was removed and stirring continued for one and a half hours at room temperature. Aldehyde (11) (828 mg, 4.12 mmol) was added via cannula in THF (10 ml + 5 ml washings), and stirring continued for a further two hours. The reaction mixture was diluted with water (30 ml) and extracted with ether (3 x 40 ml). The combined organic extracts were washed with brine (60 ml), dried with magnesium sulfate and concentrated in vacuo. The crude oil was purified by flash column chromatography (5% ether in pet.ether) to afford the required enol ether (12) as a 40:60 mixture of *cis* and *trans* isomers, (not separated) (873 mg, 92%). The purity of this mixture was checked by 200 MHz nmr and the mixture taken on and used in the following step without further characterisation.

¹H NMR (200 MHz, CDCl₃) δ: 1.52 (2.4H, s, CCH₃(CH₃)), 1.62 (3.6H, s, CCH₃CH₃), 3.49 (1.8H, s, = OCH₃), 3.53 (1.2H, s, OCH₃), 3.73 (1.2H, s, NCH₃), 3.74 (1.6H, s, NCH₃), 4.60 (0.4H, d, J = 6.9 Hz, =CHOMe), 5.13 (0.6H, d, J=12.7 Hz, =CHOMe), 5.78 (0.4H, d, J=6.9 Hz, CH=CHOMe), 6.32 (0.6H, d, J=12.7 Hz, CH=CHOMe), 6.83 (1H, s, =CHNH), 7.02-7.40 (3H, m, 3 x ArH), 7.73-7.78 (1H, m, ArH).

(g) Preparation of Aldehyde (13)

To a stirred solution of enol ether (12) (854 mg, 3.73 mmol) in dioxane (40 ml) and water (10 ml) at room temperature was added p-toluenesulfonic acid monohydrate (100 mg, 0.526 mmol), the resulting mixture was heated to

- 40 -

60°C for sixteen hours. The reaction mixture was then diluted with water (40 ml) and extracted with ether (3 x 50 ml), the combined organic extracts were washed with brine (75 ml), dried with magnesium sulfate and concentrated in vacuo. The crude oil was purified by flash column chromatography (20% ether in pet.ether) to afford the desired aldehyde (13) as an off-white solid (696.2 mg, 86%).

10 Mp: 39-40°C

IR (CHCl₃ soln): 1718 (s, C=O), 1615, 1546 (w, C=C).

¹H NMR (400 MHz, CDCl₃) δ: 1.55 (6H, s, CCH₃(CH₃)), 2.83 (2H, d, J=3.1 Hz, CH₂CHO), 3.74 (3H, s, =NCH₃), 6.82 (1H, s, =CHNH), 7.10 (1H, dt, J=1.0, 7.3 Hz ArH), 7.24 (1H, dt, J=1.0, 7.3 Hz, ArH), 7.32 (1H, dd, J=1.0, 7.3 Hz, ArH), 7.56 (1H, dd, J=1.0, 7.3 Hz ArH), 9.51 (1H, t, J=3.1 Hz CHO).

20 ¹³C NMR (75.3 MHz, CDCl₃) δ: 204.1, 137.9, 125.6, 125.3, 121.4, 121.3, 120.7, 118.7, 109.6, 54.7, 33.6, 32.6, 29.2.

LRMS (EI): 215 (M⁺, 45%), 172 (100%).

25 HRMS (EI) Calcd. for C₁₄H₁₇ON: 215.13101; found: 215.13103.

Anal. Calcd for : C₁₄H₁₇ON: C, 78.10; H, 7.96; N, 6.51.
Found: C, 78.22; H, 7.98; N, 6.41.

30 (h) Preparation of Acid (14)

To a solution of aldehyde (13) 234 mg, 1.09 mmol) in tert-butyl alcohol (6 ml) at room temperature was added 2-methyl-2-butene (8.0 ml, 2.0M in THF, 16.3 mmol). To the resulting solution was added a solution of sodium chlorite

- 41 -

(148 mg, 1.63 mmol) and sodium hydrogen phosphite (600 mg, 4.36 mmol) in water (6 ml). The resulting solution was stirred for twenty minutes at room temperature and then diluted with water (10 ml), acidified to pH 1-2 with dilute hydrochloric acid and extracted with ethyl acetate (3 x 25 ml). The combined organic extracts were concentrated in vacuo with trace amounts of water being removed by azeotropic distillation with toluene. The resulting crude mixture was purified by flash column chromatography (50% ether in pet.ether + 1% acetic acid) to afford the acid (14) as an off white solid.

Mp: 139-140°C

IR (CHCl₃ soln): 3054, 2981 (s, br, OH), 1705 (s, C=O), 1620, 1540 (w, C=C).

¹H NMR (400 MHz, CDCl₃) δ: 1.63 (6H, s, CCH₃(CH₃)), 2.88 (2H, c, CH₂CO₂H), 3.74 (3H, s, = NCH₃), 6.87 (1H, s, = CHNH), 7.10 (1H, dt, J=8.0 Hz ArH), 7.24 (1H, t, J=8.0 Hz, ArH), 7.32 (1H, d, J=8.0 Hz, ArH), 7.56 (1H, d, J= 8.0 Hz ArH).

¹³C NMR (75.3 MHz, CDCl₃) δ: 178.4, 137.7, 125.7, 125.1, 122.4, 121.2, 120.7, 118.5, 109.5, 46.8, 34.0, 32.5, 28.3.

LRMS (EI): 231 (M⁺, 23%), 216 (7%), 172 (100%).

HRMS (EI) Calcd. for C₁₄H₁₇O₂N: 231.12593; found: 231.12586.

(i) Preparation of Auxiliary (15)

To a stirred cooled (-78°C) solution of acid (14) (269 mg, 1.16 mmol) in THF (20 ml) under argon was added triethylamine (243 µl, 1.75 mmol) and then trimethylacetyl

- 42 -

chloride (158 μ l, 1.28 mmol), the resulting mixture was warmed to 0°C, stirred for one hour and then re-cooled to -78°C. In a second flask butyllithium (1.27 ml, 1.53M in hexanes, 1.93 mmol) was added drop wise to a stirred
5 cooled (-78°C) solution of (4S)-(-)-4-isopropyl-2-oxazolidinone (250 mg, 1.94 mmol) in THF (8 ml) under an argon atmosphere, and the resulting lithiated oxazolidinone was added via cannula to the reaction flask. Stirring was continued for one hour and then water (20 ml)
10 was added and the reaction mixture was allowed to warm to room temperature whereupon it was extracted with ether (3 x 30 ml). The combined organic extracts were washed with brine (30 ml) and dried over magnesium sulfate, and concentrates in vacuo. The crude yellow oil was purified
15 by flash column chromatography (40% ether in pet.ether) to afford the desired compound (15) as an off white solid (313 mg, 78%).

IR (CHCl₃ soln): 1777, 1693 (s, C=O), 1615, 1540 (w, C=C).

20

¹H NMR (400 MHz, CDCl₃) δ : 0.67 (3H, d, J=6.9 Hz, CH(CH₃)CH₃), 0.77 (3H, d, J=6.9 Hz, CH(CH₃)CH₃), 1.59 (3H, s, CCH₃(CH₃)), 1.61 (3H, s, CCH₃(CH₃)), 2.14 (1H, m, CH(CH₃)CH₃), 3.48 (2H, s, CH₂CON), 3.71 (3H, s, NCH₃), 3.71 (1H, br, t, J= 9.0 Hz, CH_AH_BO), 3.97 (1H, dd, J=9.0, 2.7 Hz CH_AH_BO), 4.18 (1H, m, CH('Pr)CH₂), 6.86 (1H, s, =CHNH), 7.07 (1H, t, J=8.0 Hz ArH), 7.16 (1H, t, J=8.0 Hz, ArH), 7.24 (1H, d, J=8.0 Hz, ArH), 7.82 (1H, d, J= 8.0 Hz ArH).

25

¹³C NMR (75.3 MHz, CDCl₃) δ : 171.5, 154.0, 137.5, 125.9, 125.6, 122.1, 121.0, 118.5, 109.3, 62.9, 58.5, 45.4, 35.0, 32.6, 29.6, 28.7, 28.5, 17.9, 14.5.

30

LRMS (EI): 231 (M⁺, 23%), 216 (7%), 172 (100%).

35

- 43 -

(j) Preparation of Azide (16)

To a stirred, cooled (-78°C) solution of oxazolidinone (15) (82.7 mg, 0.242 mmol) in THF (3 ml) under an argon atmosphere was added potassium bis(trimethylsilyl) amide (0.73 ml, 0.4M in THF, 0.29 mmol) and the resulting solution was stirred at -78°C for one hour. A solution of triisopropylsulfonyl azide (97 mg, 0.315 mmol) in THF (2 ml) pre-cooled to -78°C was added via cannula and after one minute the reaction was quenched by addition of glacial acetic acid (100 ml), warmed to 40°C (water bath) and stirred for a further hour. The reaction mixture was then partitioned between dichloromethane (10 ml) and dilute brine (10 ml), and the layers separated. The aqueous phase was extracted with dichloromethane (2 x 10 ml) and the combined organic extracts washed with sodium hydrogen carbonate (10 ml, sat.aq.), brine (10 ml), dried with magnesium sulfate and concentrated *in vacuo*. The resulting crude oil was purified by flash column chromatography (30% ether in pet.ether) to afford a mixture of the desired compound (16) and trisopropylsulphonylamine (56.7 mg total, inseparable, estimated 45 mg of desired compound, approx. 50%).

¹H NMR (200 MHz, CDCl₃) δ: 0.70 (3H,d,J= 6.9 Hz, CH(CH₃)CH₃), 0.76 (3H, d,J=6.9 Hz, CH(CH₃)CH₃), 1.62 (3H,s,CCH₃(CH₃)), 1.66 (3H,s, CCH₃(CH₃)), 2.16 (1H,m, CH(CH₃)CH₃) 3.66-3.73 (6H,m,CH₂CON, NCH₃, CH('Pr)CH₃), 5.66 (1H,s,CHN₃), 6.94 (1H,s,=CHNH), 7.06 (1H,t,J=8.0 Hz, ArH), 7.16 (1H,t,J=8.0 Hz,ArH), 7.26 (1H,d,J=8.0 Hz,ArH), 7.75 (1H,d,J=8.0 Hz, ArH).

(k) Preparation of Boc-auxiliary (17)

- 44 -

A mixture of azide (16) (58 mg, semi-crude, <0.152 mmol) 10% palladium on charcoal (30 mg), and di-tert-butyl dicarbonate (66mg, 0.304 mmol) in ethyl acetate (4 ml) was flushed with argon and then hydrogen and stirred under a hydrogen balloon for sixteen hours at room temperature. The palladium was removed via filtration through celite, the solvent removed in vacuo and the crude mixture purified by flash column chromatography (50% ether in pet.ether) to afford the desired compound (17) (21.8 mg, <50%).

^1H NMR (200 MHz, CDCl_3) δ : 0.71 (3H,d,J=6.9 Hz, $\text{CH}(\text{CH}_3)\text{CH}_3$), 0.73 (3H,d, J=6.9 Hz, $\text{CH}(\text{CH}_3)\text{CH}_3$), 1.40 (9H,s, $\text{C}(\text{CH}_3)_3$), 1.53 (3H,s, $\text{CCH}_3(\text{CH}_3)$), 1.59 (3H,s. $\text{CCH}_3(\text{CH}_3)$), 2.10 (1H,m, $\text{CH}(\text{CH}_3)\text{CH}_3$), 2.58 (1H,m, $\text{CH}_2\text{H}_2\text{O}$), 3.66-3.73 (5H,m, $\text{CH}_2\text{H}_2\text{O}$, NCH_3 , $\text{CH}(\text{iPr})\text{CH}_2$), 5.28 (1H, br, NH), 6.03 (1H, br,d, CHNHBOc), 6.98 (1H,s, $=\text{CHNH}$), 7.02 (1H,t,J=8.0 Hz, ArH), 7.16 (1H,t,J=8.0 Hz, ArH), 7.24 (1H,d,J=8.0 Hz, ArH), 7.72 (1H,d,J=8.0 Hz, ArH).

(1) Preparation of Methyl ester of (17)

To a solution of (17) (13 mg, 0.0285 mmol) in THF (1 ml) and water (0.3 ml) at 0°C was added lithium hydroxide (8 mg, excess) as a solid in one portion. The resulting suspension was stirred at room temperature for sixteen hours, and then acidified with 1N hydrochloric acid, and the solvent removed in vacuo. The resulting crude white solid was suspended in ether and an ethereal solution of diazomethane added, until the yellow colour of the diazomethane persisted in the reaction mixture and tlc analysis showed complete consumption of starting material. The excess diazomethane was removed under a stream of argon and the remaining solvent removed in vacuo. The resulting crude oil was purified by flash column

- 45 -

chromatography (40% ether in pet.ether) to afford the desired compound methyl ester of (17) (7.7 mg. 75%).

¹H NMR (400 MHz, CDCl₃) δ: 1.38 (9H,s,C(CH₃)₃), 1.47
5 (3H,s,CCH₃(CH₃)), 1.52 (3H,s,CCH₃(CH₃)), 3.45 (3H,s, CO₂CH₃),
3.72 (3H,s,NCH₃), 4.70 (1H,d,br,NH), 5.05
(1H,d,br,CHNH₂Boc), 6.81 (1H,s,=CHNH), 7.07 (1H,t,J=8.0 Hz
ArH), 7.21 (1H,t,J=8.0 Hz, ArH), 7.30 (1H,d,J=8.0 Hz,ArH),
7.80 (1H,d,J= 8.0 Hz ArH).

10

(m) Alternative route to methyl ester of (17)

Reversal of the above two steps allows for the synthesis of the same compound in comparable yield.

15

(n) Preparation of methylated methyl ester

To a suspension of sodium hydride (excess) and methyl iodide (excess) in dry THF (0.5 ml) under an argon
20 atmosphere was added the methyl ester of (17) (10.4 mg
0.0289 mmol) in THF (1 ml) via cannula and the resulting
mixture was stirred for sixteen hours at room temperature.
Water was added and the mixture was extracted with ether,
the combined organic extracts were washed with brine,
25 dried with magnesium sulfate and concentrated in vacuo.
The crude oil was purified by flash column chromatography
(40% ether in pet.ether) to afford the desired N-
methylated methyl ester as a colourless oil (3.1 mg,
approx. 30%).

30

¹H NMR (400 MHz, CDCl₃) δ: 1.42 (9H,s,C(CH₃)₃), 1.52
(3H,s,CCH₃(CH₃)), 1.64 (3H,s,CCH₃(CH₃)), 2.70 (3H,s,(split)
NCH₃Boc), 3.45 (3H,s, split, CO₂CH₃), 3.71 (3H,s,NCH₃), 5.40
(1H,s,split, CHNCH₃Boc), 6.98 (1H,s,= CHNH), 7.00-7.25
35 (3H,m, ArH), 7.72 (1H,d,br,split,ArH).

- 46 -

(o) Preparation of (18)

The procedure of step (1) was followed, but the esterification with diazomethane was omitted.

5

(p) Preparation of (19)

N-Boc-amine acid (18) (10 mmol) and methyl iodide (5 ml; 80 mmol) were dissolved in THF (30 ml) and the solution was cooled to 0°C under argon. Sodium hydride dispersion (1.32 g; 30 mmol) was added cautiously with gentle stirring. The suspension was stirred at room temperature for 16 h. Ethyl acetate (50 ml) was then added (to consume the sodium hydroxide formed from the excess sodium hydride), followed by water, drop wise, to destroy the excess sodium hydride. The solution was evaporated to dryness, and the oily residue partitioned between ether (30 ml) and water (100 ml). The ether layer was washed with 5% aqueous NaHCO₃ (50 ml) and the combined aqueous extracts and acidified to pH 3 with aqueous citric acid. The product was extracted into ethyl acetate (3 x 25 ml), the extract washed with water (2 x 25 ml), 5% aqueous sodium thiosulphate (2 x 25 ml; to remove iodine), and water (2 x 25 ml), dried over MgSO₄ and evaporated to give a pale yellow oil of (19).

3. Coupling of amino acids

N-Boc amino acid (19) (1 mmol), amino acid methyl ester of moiety B (1.1 mmol) and coupling agent py-BOP (1.1 mmol) were dissolved in CH₂Cl₂ (10 ml) under argon. TIEA (3 mmol) was added and the reaction was stirred for 1 h at room temperature. Excess solvents were removed in vacuo yielding a yellow oily residue which was redissolved in EtOAc (50 ml). Washing the EtOAc solution with 10%


- 47 -

citric acid (2 x 25 ml), water (25 ml), 5% NaHCO₃ (2 x 25 ml), water (25 ml), following by drying over anhydrous MgSO₄ and normal phase silica gel chromatography afforded the protected peptide A-B as a white amorphous solid.

5

Coupling of the peptide A-B to amino acid moiety of C (5) was carried out in a similar way.


- 48 -

Scheme 1**Synthesis of N-Me-homo vinyllogous valine ethyl ester (MHVV-OEt)**

SUBSTITUTE SHEET (RULE 26)

- 49 -


Scheme 2



SUBSTITUTE SHEET (RULE 26)

- 50 -

Scheme 2 (cont'd)



SUBSTITUTE SHEET (RULE 26)





- 51 -

Testing of compounds

1. The cytotoxicity of compounds described herein have been tested as described in J. Immunol. Methods, 65, 55-63, 1983 and the results are provided in Table 4 below, wherein P388 refers to in vitro test versus murine leukaemia P388, U373 refers to in vitro human glioblastoma/astrocytoma U373, HEY refers to in vitro human ovarian carcinoma HEY, MCF7 refers to in vitro human breast cancer MCF7.
2. In in vivo tests as described in NIH Publication No. 84-2635, "In Vivo Cancer Models", Developmental Therapeutic Program, Division of Cancer Treatment, National Cancer Institute, Bethesda, Maryland, hemiasterlin has been found to be cytotoxic. In mice injected with 1×10^6 P388 leukaemia cells, hemiasterlin resulted in a %TC of 223 after 5 daily doses of 0.45 μ g begun 24 hours after implantation. There were several long term survivors in the experiment.




- 52 -

TABLE 4

compound	IC ₅₀ Values (µg/ml)				
	P388	U373	HEY	MCF7	cell mitosis
hemiasterlin (Compound A) 	4.57 x 10 ⁻⁵	1.2 x 10 ⁻²	1.4 x 10 ⁻³	1.58 x 10 ⁻⁴	1.58 x 10 ⁻⁴
hemiasterlin-A (Compound 2) 	1.71 x 10 ⁻⁶	1.5 x 10 ⁻³	7.6 x 10 ⁻³	1.54 x 10 ⁻³	1.02 x 10 ⁻³
hemiasterlin-B (Compound 3) 	7.0 x 10 ⁻³		1.6 x 10 ⁻²	1.50 x 10 ⁻²	9.96 x 10 ⁻³
criamide-B 	73 x 10 ⁻³	0.27	0.19	6.8	

- 53 -

TABLE 4 (Continued)

	IC ₅₀ Values (µg/ml)				
Geodiamolide G (compound)		7.7	8.6		
hemiasterlin-OMe 				5.4 x 10 ⁻³	2.16 x 10 ⁻⁴
dihydrohemiasterlin 					1.06 x 10 ⁻³
 Totally Synthetic analog				0.1	

SUBSTITUTE SHEET (RULE 26)

- 54. -

3. Compounds described herein were comparatively tested for their antimitotic activity against human mammary carcinoma MCF7 cells.

5 MCF7 cells were grown as a monolayer in RPMI supplemented with 15% fetal calf serum and antibiotics at 37°C in humidified 10% CO₂. All compounds were dissolved in dimethyl sulfoxide except for vinblastine (a known drug) which was a 1mg/ml solution in physiological saline.
10 Exponentially growing MCF7 cells were treated with different drug concentrations for 20 h, prepared for chromosome spreads, and the percentage of mitotic cells determined by fluorescence microscopy. The results are shown in Figures 1 and 2. Hemiasterlin, Hemiasterlin A and
15 modified compounds were very potent antimitotic agents, with IC₅₀ values of 0.3 nM and 3 nM respectively. Hemiasterlin and Hemiasterlin A were more potent than Taxol, Vinblastine and Nocodazole (all known drugs).

20 The effect of Hemiasterlin and Hemiasterlin A on the morphology of their mitotic spindles was examined by indirect immunofluorescence using a monoclonal antibody to β -tubulin and the distribution of their chromosomes using the fluorescent DNA dye bisbenzimidazole. In the presence of
25 hemiasterlin A at 2 nM no completely normal spindles were seen. Some cells showed relatively minor abnormalities in which a bipolar spindle was present but the astral microtubules were considerably longer than normal and the chromosomes were not completely confined to the metaphase
30 plate. Most commonly cells had multiple asters, and the chromosomes were distributed in a spherical mass. Half-maximal concentrations of taxol, vinblastine and nocodazole produced the same types of abnormal spindle as hemiasterlin A. Hemiasterlin A at 10 nM, the lowest
35 concentration causing maximal mitotic arrest in MCF7

- 55 -


cells, caused microtubule depolymerisation in mitotic cells. This was also the case for high concentrations of vinblastine and nocodazole. Taxol at high concentrations had a quite different effect, causing bundling of
5 cytoplasmic microtubules in interphase cells and very dense multiple asters in mitotic cells.

These results show that Hemiasterlins cause mitotic arrest and produce abnormal mitotic spindles. They can be
10 used in lieu of other antimitotic drugs in procedures that require blocking cells in mitosis, such as the preparation of mitotic spreads for karyotype analysis. They can also be used to probe microtubule function in mitotic cells.

- 56 -

CLAIMS

1. The use of a hemiasterlin compound of general formula



wherein:

- R_1 and R_{70} independently represent a hydrogen atom or an optionally substituted alkyl or acyl group;
- 15 R_2 represents a hydrogen atom or an optionally substituted alkyl or acyl group or is absent when R_6 represents a group $-CH=$ as hereinafter described;
- R_{73} represents a hydrogen atom or an optional substituent or is absent when R_6 represents a methylene group or a group $-CH=$ as hereinafter described;
- 20 Y represents an optional substituent;
- n represents 0, 1, 2, 3, or 4;
- R_3 represents a hydrogen atom, or an optionally substituted alkyl group;
- 25 R^{74} represents a hydrogen atom, a hydroxy group or an optionally substituted alkyl or acyl group;
- R_7 represents a hydrogen atom or an alkyl group;
- R_{75} represents an optionally substituted alkyl group; and
- i) R_6 and R_{71} independently represent a hydrogen atom or
- 30 an optionally substituted alkyl or acyl group; and
- R_{72} represents a hydrogen atom;
- or
- ii) R_{71} represents a hydrogen atom or an optionally substituted alkyl or acyl group and R_{72} represents a
- 35 hydrogen atom or R_{71} and R_{72} each represent radicals so that

- 57 -

a double bond is formed between the carbon atoms to which they are attached; and

5 R_4 represents an optionally substituted methylene group bonded to the indole moiety thereby to form a tricyclic moiety; or

R_4 represents an optionally substituted group $-CH=$ bonded to the indole moiety thereby to form an aromatic tricyclic moiety;

10 for the manufacture of a medicament for use in therapy.

2. Use of a hemiasterlin according to Claim 1, wherein R_2 represents a hydrogen atom or an acyl group.


15 3. Use of a hemiasterlin according to Claim 1 or Claim 2, wherein n represents 0.

20 4. Use of a hemiasterline according to any preceding claim, wherein R_{73} represents a hydrogen atom.

5. Use of a hemiasterline according to any preceding claim, wherein R_6 represents an alkyl group.

25 6. Use of a hemiasterline according to any preceding claim, wherein R_{71} represents a hydrogen atom or an optionally substituted alkyl or acyl group.

30 7. Use of a hemiasterline according to any preceding claim, wherein R_{73} represents a group of general formula



II

- 58 -

wherein Q represents an optionally substituted -CH₂-, -CH₂CH₂-, -CH₂CH₂CH₂-, -CH₂CHCH-, -CH₂C.C.- or phenylene moiety; and

- 5 X represents a group -OR₈, -SR₈, or -NR₉R₁₀ wherein R₈, R₉ and R₁₀ independently represent a hydrogen atom or an optionally substituted alkyl group.
8. Use of a hemiasterlin according to Claim 7, wherein
10 X represents a group -OR₈ wherein R₈ represents a hydrogen atom or a methyl group.
9. Use of a hemiasterline according to Claim 7 or Claim 8, wherein X represents a group -NR₉R₁₀ wherein R₉
15 represents a hydrogen atom or an alkyl group, and R₁₀ represents a substituted alkyl group.
10. Use of a compound of general formula I as described
20 in any of Claims 1 to 9, as an antimitotic compound.
11. A novel compound of general formula I as described in any of Claims 1 to 9, but excluding a single compound of general formula I wherein R₁ represents methyl, R₂ represents a hydrogen, R₇₀ represents methyl, R₇₁ represents
25 methyl, R₇₃ represents hydrogen, n represents 0, R₃ represents t-butyl, R₇₄ represents hydrogen, R₆ represents methyl, R₇ represents methyl, R₇₂ represents hydrogen and R₇₅ represents -CH(CH(CH₃)₂)CH.CCH₃.COOH.
12. A hemiasterlin of general formula I described in
30 Claim 1, wherein
R₁ represents a hydrogen atom;
R₂ represents a hydrogen atom, or an alkyl group, or an acyl group;

- 59 -

- R_3 represents a hydrogen atom, or an optionally substituted alkyl group;
 n represents 0;
 R_{70} and R_{71} independently represent a hydrogen atom or optionally substituted alkyl group, but preferably each represent a methyl group;
 R_{72} , R_{73} and R_{74} represent hydrogen atoms;
 R_7 represents a hydrogen atom or an alkyl group;
 R_6 represents a hydrogen atom, or an optionally substituted alkyl group, or a methylene group bonded to the indole moiety thereby to form a tricyclic moiety;
 R_{75} represents a group of general formula III described above wherein R_4 represents a hydrogen atom, or an optionally substituted alkyl group; R_5 represents a hydrogen atom or an alkyl group; R_{76} and R_{77} represent radicals as described; and X represents a group $-OR_8$ or a group $-NR_9R_{10}$, wherein R_8 , R_9 and R_{10} independently represent a hydrogen atom or an optionally substituted alkyl group.
13. A hemiasterlin of general formula I described in Claim 1, wherein
 R_1 represents a hydrogen atom or an alkyl group;
 R_2 represents an acyl group;
 R_3 represents a hydrogen atom, or an optionally substituted alkyl group;
 n represents 0;
 R_{70} and R_{71} independently represent a hydrogen atom or optionally substituted alkyl group, but preferably each represent a methyl group;
 R_{72} , R_{73} and R_{74} represent hydrogen atoms;
 R_7 represents a hydrogen atom or an alkyl group;
 R_6 represents a hydrogen atom, or an optionally substituted alkyl group, or a methylene group bonded to the indole moiety thereby to form a tricyclic moiety;

- 60 -


R_{75} represents a group of general formula III described above wherein R_4 represents a hydrogen atom, or an optionally substituted alkyl group; R_5 represents a hydrogen atom or an alkyl group; R_{76} and R_{77} represent radicals as described; and X represents a group $-OR_8$ or a group $-NR_9R_{10}$, wherein R_8 , R_9 and R_{10} independently represent a hydrogen atom or an optionally substituted alkyl group.

14. A criamide of general formula I described in Claim 1, wherein

R_1 represents a hydrogen atom or an alkyl group;
 R_2 represents a hydrogen atom, or an alkyl group, or an acyl group;
 R_3 represents a hydrogen atom, or an optionally substituted alkyl group;
 n represents 0;
 R_{70} and R_{71} independently represent a hydrogen atom or optionally substituted alkyl group, but preferably each represent a methyl group;
 R_{72} , R_{73} and R_{74} represent hydrogen atoms;
 R_6 represents a hydrogen atom, or an optionally substituted alkyl group, or a methylene group bonded to the indole moiety thereby to form a tricyclic moiety;
 R_{75} represents a group of general formula III described above wherein R_4 represents a hydrogen atom, or an optionally substituted alkyl group; R_5 represents a hydrogen atom or an alkyl group; R_{76} and R_{77} represent radicals as described; and X represents a group $-NR_9R_{10}$, wherein R_9 and R_{10} independently represent a hydrogen atom or an optionally substituted alkyl group.

15. A geodiamolide compound of general formula

- 61 -



wherein:

R₅₁ represents an alkyl group;

R₅₂ represents a hydrogen atom or an alkyl group; and

15 A represents a halogen atom.


20

25


30

35

1/1

Figure 1.

Comparison of antimittotic activity of Hemiasterlins with that of known antimittotic agents.

Figure 2.

Antimittotic activity of chemically modified Hemiasterlin.

INTERNATIONAL SEARCH REPORT

Inter nal Application No

PC1/GB 96/00942

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07K5/027 C07K5/078 C07K5/083

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	TETRAHEDRON (1995), 51(39), 10653-62 CODEN: TETRAB;ISSN: 0040-4020, 25 September 1995, XP002012447 COLEMAN, JOHN E. ET AL: "Cytotoxic peptides from the marine sponge Cymbastela sp." see the whole document ---	1-15
X	TETRAHEDRON LETT. (1994), 35(25), 4453-6 CODEN: TELEAY;ISSN: 0040-4039, 1994, XP002012448 TALPIR, R. ET AL: "Hemiasterlin and geodiamolide TA;two new cytotoxic peptides from the marine sponge Hemiasterella minor (Kirkpatrick)" compounds 2-3 see page 4455, paragraph 2 --- -/--	1-13,15

☒ Further documents are listed in the continuation of box C.☐ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

3 September 1996

Date of mailing of the international search report

13.09.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Fuhr, C

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 96/00942

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JOURNAL OF ORGANIC CHEMISTRY, vol. 59, no. 11, 3 June 1994, EASTON US, pages 2932-2934, XP002012449 P. CREWS ET AL.: "Milnamide A, an Unusual Cytotoxic Tripeptide from the Marine Sponge Auletta cf. constricta" compounds 1-3 see page 2932, left-hand column, paragraph 2	1-13
A	--- JOURNAL OF ORGANIC CHEMISTRY, vol. 52, no. 14, 10 July 1987, EASTON US, pages 3091-3093, XP002012450 W.R. CHAN ET AL.: "Stereostructures of Geodiamolides A and B, Novel Cyclodepsipeptides from the Marine Sponge Geodia sp." compounds 1-3 see page 3092, right-hand column, paragraph 5	15
P,A	--- EXPERIMENTAL HEMATOLOGY, vol. 23, no. 7, July 1995, pages 583-587, XP000601558 I. FABIAN ET AL.: "Growth modulation and differentiation of acute myeloid leukemia cells by jaspamide" see the whole document -----	1-15